

LETTER TO THE EDITOR

Characterization of damage in Portuguese lupus patients

Sir,

We read with interest the report 'Characterization of damage in Portuguese lupus patients: analysis of a national lupus registry'¹ and wish to emphasize that the frequency and cumulative damage score present in a lupus patient population is also determined by the type of organ damage.

In the registry study of 976 patients by Gonçalves et al.,¹ the average SDI score was 0.71 ± 1.22 and the frequency of damage was 37.4% after a 17 ± 9.4 year follow-up. This was considered by the authors to be lower in Portuguese patients when compared to other series, namely the University College London Hospital (UCLH) cohort which reported 50% damage at 10 year follow-up² and the SLICC cohort, which observed 51% damage at six years follow-up.³ Of note, in the UCLH cohort, when measured between 15 and 20 years of follow-up, most of the damage accrual occurred in the musculoskeletal system.

A recent retrospective study of our single-centre hospital-based cohort consisting of 98 Portuguese lupus patients, in which every patient was clinically evaluated at the time of study completion, revealed a similar frequency of accrued damage equally measured by SDI values ≥ 1 .⁴ In comparison, although our cohort comprised a similar frequency of articular disease (approximately 70%), there was a higher frequency of neuropsychiatric (NP) disease (22%) than in the national register (11%). In addition, in our cohort, the patients with damage were enriched for NP disease ($n = 16/22$, 72%) versus non-NP patients ($n = 17/76$, 22%). Our average SDI scores were higher: 2.31 ± 1.25 for the NP and 1.53 ± 0.80 for the non-NP group. This difference was statistically significant ($P = 0$, 0001) and was largely due to damage of the neurologic system that occurred in 14 of the 16 NP patients. Both NP and non-NP patients had a similar mean age and disease

duration (age: 44 ± 12 and 45 ± 14 years, disease duration: 13 ± 7 and 13 ± 8 years, respectively). SDI scores increased with the duration of the disease in both groups, and were initially 0.44 ± 0.89 for NP and 0 for non-NP patients.

It is well conceivable that Lupus patients with a predominant articular phenotype should have low damage accrual after a long follow-up. However, this is a very heterogeneous disease, and this subset of patients is not representative of the overall population of lupus patients, when other clinical phenotypes are taken into consideration.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors have no conflicts of interest to declare.

References

- 1 Gonçalves MJ, Sousa S, Inês LS, et al. Characterization of damage in Portuguese lupus patients: analysis of a national lupus registry. *Lupus* 2015; 24: 256–262.
- 2 Chambers SA, Allen E, Rahman A, et al. Damage and mortality in a group of British patients with systemic lupus erythematosus followed up for over 10 years. *Rheumatology (Oxford)* 2009; 48: 673–675.
- 3 Bruce IN, O'Keefe AG, Farewell V, et al. Factors associated with damage accrual in patients with systemic lupus erythematosus: results from the Systemic Lupus International Collaborating Clinics (SLICC) Inception Cohort. *Ann Rheum Dis*. Epub: May 2014 DOI: 10.1136/annrheumdis-2013-205171.
- 4 Moraes-Fontes MF, Lúcio I, Santos C, et al. Neuropsychiatric features of a cohort of patients with systemic lupus erythematosus. *ISRN Rheumatology*. Epub Nov 2012 DOI: 10.5402/2012/989218.

MF Moraes-Fontes and N Riso
Hospital Curry Cabral, Lisboa, Portugal

Correspondence to: Maria Francisca Moraes-Fontes, Hospital Curry Cabral, Centro Hospitalar Lisboa Central Rua da Benficiência Lisboa, 1069-166 Portugal.

Email: mffontes@igc.gulbenkian.pt

Received 4 February 2015; accepted 17 March 2015

Copyright of *Lupus* is the property of Sage Publications, Ltd. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.